

Types of Inoculants & things to consider when using them





PEAT

Description – Finely ground sterilised peat containing rhizobia with adhesive.

Storage – Refrigerate at 4°C. Do not freeze.

Application – Create peat slurry by mixing with clean water. Seeds can be either coated before sowing or peat can be placed into porous bag and added to water for liquid injection. Sow within 5hrs.



LIQUID

Description – Suspension of rhizobia in protective liquid formulation with adhesive 'sticker' powder.

Storage – Refrigerate at 4°C. Do not freeze.

Application – Mix inoculant and adhesive with clean water and either apply directly to seed or add into inoculation tank for liquid injection. Sow within 5hrs of application.



FREEZE-DRIED

Description – Vial of soluble powder with additional protective agent. Activated when reconstituted with liquid.

Storage – Refrigerate at 4°C. Do not freeze.

Application – Mix freeze-dried powder with water and either apply directly to seed or add to inoculation tank for liquid injection. Sow within 5hrs.

- Inoculants contain living bacteria. Protect them by storing and applying them according to the packet.
- Select the correct inoculant group for the type of legume.
- Apply inoculant and sow in the late evening to avoid high day time temperatures.
- Consider doubling the rate of inoculant for adverse conditions.



• Look for the Australian Inoculants Research Group (AIRG) Green Tick Logo on your inoculant for a quality guaranteed product.

• Take care with using clean, potable water when preparing suspensions and ensure your equipment and tanks are clean.

• Speak to a HCPSL agronomist for more inoculant information. You can also view useful factsheets using the QR code.

Legume Inoculant Selector

Selecting the right inoculant group for each legume type is critical for ensuring you get nodulation and good nitrogen fixation.

Inoculant groups are determined by the strain of rhizobia that is specifically suited to a type of legume host.

INOCULANT GROUP	LEGUME HOST PLANT
Н	Soybean
I	Cowpea Mungbean
J	Pigeon Pea Lablab
Μ	Sunn Hemp* Velvet bean
N	Chickpea
Р	Peanut Groundnut
Special (CB1717 strain)	Burgundy bean

*If M strain inoculant is unavailable, group I is suitable for use with Sunn Hemp.

If you are working with mixed legumes, remember to supply inoculant at a rate appropriate for the whole mix to ensure sufficient coverage.

Leftover inoculant can be stored for later use. Aim to remove air from the packet before sealing and storing refrigerated at 4°C. Make sure to use the inoculant before the labelled expiry date.



Project Catalyst is funded by the partnership between the Australia Government's Reef Trust and the Great Barrier Reef Foundation, The Coca-Cola Foundation and WWF-Australia.

Disclaimer: Copyright 2021. All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise, without the prior permission of HCPSL. Disclaimer In this disclaimer a reference to "we", "us" or "our" means Herbert Cane Productivity Services Ltd (HCPSL) and our Directors, Officers, Agents and Employees. Although we do our best to present information that is correct and accurate, we make no warranties, guarantees or representations about the suitability, reliability, currency or accuracy of the information we present in this report for any purposes. Subject to any terms implied by law and which cannot be excluded, we accept no responsibility for any loss, damage, cost or expense incurred by you as a result of the use of, or reliance on, any materials and information appearing in this report and you agree that we will not be liable for any loss or damage whatsoever (including through negligence) arising out of, or in connection with the use of this report. We recommend that you contact our staff before acting on any information provided in this report.